


Alexa Fluor® 488 anti-human CD11a

Catalog # / Size:	2106080 / 100 tests
Clone:	HI111
Isotype:	Mouse IgG1, κ
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 488 under optimal conditions.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Workshop Number:	IV N231
Concentration:	NULL



Human peripheral blood lymphocytes stained with HI111 Alexa Fluor® 488

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.

Application Notes: Clone HI111 epitope maps to the top region of the I domain that is close to the putative ligand-binding site surrounding the MIDAS (metal ion-dependent adhesion site). HI111 is specific for the closed confirmation of the integrin.⁸ Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections, Western blotting², and blocking of cell-cell interaction and inhibition the binding of ICAM-14. This clone was tested in-house and does not work on formalin fixed paraffin-embedded (FFPE) tissue. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 301214).

Application References:

1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press New York.
2. Leite F, *et al.* 2002. *Infect. Immun.* 70:4336.
3. Jiang Y, *et al.* 2005. *Clin. Hemorheol. Microcircul.* 32:261.
4. Béchard D, *et al.* 2001. *J. Immunol.* 167:3099.
5. Sithu SD, *et al.* 2007. *J. Biol. Chem.* doi:10.1074/jbc.M611273200.
6. Choi EY, *et al.* 2008. *Blood* 111:3607. [PubMed](#)
7. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
8. Ma Q, *et al.* 2002. *J. Biol. Chem.* 277:10638.

Description: CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1 α chain and integrin α_L subunit. CD11a non-covalently associates with integrin β_2 (CD18) to form LFA-1. It is expressed on all leukocytes, including B and T

lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

- Antigen** 1. Lub M, *et al.* 1995. *Immunol. Today* 16:479.
References: 2. Parsons J. 1996. *Curr. Opin. Cell Biol.* 8:146.