

APC/Fire™ 750 anti-human CD11b (activated)

Catalog # / Size: 2107100 / 100 tests
2107095 / 25 tests

Clone: CBRM1/5

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD01.01

Concentration: Lot-specific



PMA-activated (10 minutes) human peripheral blood granulocytes were stained with CD11b (clone CBRM1/5) APC/Fire™ 750 (filled histogram) or mouse IgG1, κ APC/Fire™ 750 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

Application Notes: The CBRM1/5 antibody recognizes a subset of CD11b molecules on neutrophils and monocytes activated with chemoattractants or phorbol-esters. This antibody does not recognize "non-activated" CD11b. The epitope recognized by CBRM1/5 is contained in the I domain of the α chain. Clone CBRM1/5 binds the I domain, close to the ligand binding site. It recognizes a conformational change of the integrin⁹.

Additional reported applications (for the relevant formats) include: immunoprecipitation, and blocking cell adhesion to fibrinogen and ICAM-1¹. The CBRM1/5 antibody binds specifically to the activated form of CD11b. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 301408).

- Application References:**
1. Diamond M, et al. 1993. *J. Cell Biol.* 120:545. (Block)
 2. Bryn T, et al. 2006. *J. Immunol.* 176:7361. [PubMed](#)
 3. Sithu SD, et al. 2007. *J. Biol. Chem.* doi:10.1074/jbc.M611273200.
 4. Barthel SR, et al. 2006. *Am J. Respir Cell Mol Biol.* 35:378. [PubMed](#)
 5. Pillay J, et al. 2010. *J. Leukocyte Biol.* 88:211. [PubMed](#)
 6. Wilson RP, et al. 2011. *Infect Immun.* 79:830. [PubMed](#)
 7. Koleva RI, et al. 2012. *Blood.* 119:4878. [PubMed](#).
 8. Eleftherious D, et al. 2012. *Neurology.* 79:2089. [PubMed](#)
 9. Oxvig C, et al. 1999. *Proc. Natl. Acad. Sci. USA* 96:2215.
 10. Muller-Edenborn B, et al. 2014. *Br J Anaesth.* [PubMed](#)
 11. Cash JL, et al. 2013. *EMBO Rep.* 14:999. (FC) [PubMed](#)

Description: The CBRM1/5 antibody reacts with an activated form of human CD11b, a 165-170 kD type I transmembrane glycoprotein also known as α_M integrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin β_2 (CD18) and is expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils as well as adhesion, phagocytosis, and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen and factor X.

Antigen 1. Stewart M, et al. 1995. *Curr. Opin. Cell Biol.* 7:690.
References: 2. Petty H, et al. 1996. *Immunol. Today* 17:209.