

Purified anti-human CD11c

Catalog # / Size: 2108005 / 25 µg
2108010 / 100 µg

Clone: 3.9

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Workshop Number: III NL707

Concentration: 0.5



Human peripheral blood granulocytes were stained with purified CD11c (clone 3.9) (filled histogram) or purified mouse IgG1, κ isotype control (open histogram), followed by anti-mouse IgG FITC.

Applications:

Applications: Flow Cytometry, Immunohistochemistry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤2.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone 3.9 preferentially binds the activated form of CD11c, is specific for the I domain of CD11c, and is able to partially block the binding of CD11c and ICAM-4. 3.9 binding is divalent cation dependent^{1,2}. While analyzing blood, it is best to use heparin as the anti-coagulant and not EDTA. Since the ability of clone 3.9 to bind to its target is divalent cation dependent, the usage of EDTA as an anti-coagulant may be detrimental to staining due to its chelating properties.

Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴, and functional assays^{5,6}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 301616). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 301632) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Knapp W, *et al.* 1989. Leucocyte Typing IV Oxford University Press. New York.
 - McMichael A, *et al.* Eds. 1987. Leucocyte Typing III Oxford University Press. New York.
 - Vainer B, *et al.* 2000. *Am. J. Surg. Pathol.* 24:1115. (IHC)
 - Ottonello L, *et al.* 1999. *Blood* 93:3505.
 - Metelitsa LS, *et al.* 2002. *Blood* 99:4166.
 - Sadhu C, *et al.* 2007. *J. Leukoc. Biol.* doi:10.1189/jlb.1106680. [PubMed](#)
 - Ihanus E, *et al.* 2007. *Blood* 109:802-810.

9. Gurer C, *et al.* 2008. *Blood* 112:1231. [PubMed](#)
 10. Asai A, *et al.* 2009. *J. Lipid Res.* 50:95. [PubMed](#)
 11. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 12. Sadhu C, *et al.* 2008. *J. Immunoass. Immunoch.* 29:42. (FC)
-

Description: CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin α_X and CR4. CD11c non-covalently associates with integrin β_2 (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing through its interactions with fibrinogen, CD54, and iC3b.

- Antigen**
References:
1. Petty H. 1996. *Immunol. Today* 17:209.
 2. Springer T. 1994. *Cell* 76:301.
 3. Ihanus E, *et al.* 2007. *Blood* 109:802-810.