## PE/Cy7 anti-human CD15 (SSEA-1)

Catalog # / 2109620 / 100 tests

Size: 2109615 / 25 tests

Clone: H198

Isotype: Mouse IgM, ĸ

Immunogen: Mouse CXCR3-transfectants Reactivity: Human, Non-human primate

Preparation: The antibody was purified by affinity

> chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.


Formulation: Phosphate-buffered solution, pH 7.2.

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Workshop Number: IV M141

Concentration: Lot-specific



Human peripheral blood granulocytes were stained with CD15 (SSEA-1) PE/Cy7 (Clone HI98) (filled histogram) or Mouse IgM

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application

Notes:

Clone HI98 has been described to specifically react with 3-fucosyl-Nacetyllactosamine (3-FAL), also called X-hapten, SSEA-1, CD15, or Lewis X. Additional reported applications (for the relevant formats) include:

immunohistochemical staining of acetone-fixed frozen sections<sup>5</sup> and

formalin-fixed paraffin-embedded tissue sections.

**Application** References:

1. Stocks SC, et al. 1990. Biochem. J. 268:275.

**Description:** 

CD15 is 3-fucosyl-N-acetyllactosamine (3-FAL), also known as Lewis X, 3-FAL, X-hapten, and SSEA-1. CD15 is expressed on granulocytes and monocytes. It has also been shown to be expressed on Langerhans cells and some malignant cells. CD15 has been implicated in adhesion, as well as chemotaxis, phagocytosis, and bactericidal activity.

Antigen References:

1. Stocks SC, et al. 1990. Biochem. J. 268:275.