

Brilliant Violet 421™ anti-human CD16

Catalog # / Size: 2110185 / 25 tests
2110190 / 100 tests

Clone: 3G8

Isotype: Mouse IgG1, κ

Immunogen: Human PMN cells


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: V NK80

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD16 (clone 3G8) Brilliant Violet 421™ (filled histogram) or mouse IgG1, κ Brilliant Violet 421™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The 3G8 antibody blocks neutrophil phagocytosis and stimulates NK cell proliferation. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁶, immunoprecipitation³, stimulation of NK cell proliferation⁴, blocking of phagocytosis⁵, and blocking of immunoglobulin binding to FcγRIII^{7,8}. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 302014). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302050) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

Application 1. Knapp W, *et al.* Eds. 1989. Leucocyte Typing IV. Oxford University Press. New

- References:** York.
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Edberg J, *et al.* 1997. *J. Immunol.* 159:3849. (IP)
 - Hoshino S, *et al.* 1991. *Blood* 78:3232. (Stim)
 - Tamm A, *et al.* 1996. *Immunol.* 157:1576. (Block)
 - Da Silva DM, *et al.* 2001. *Int. Immunol.* 13:633. (IHC)
 - Holl V, *et al.* 2004. *J. Immunol.* 173:6274. (Block)
 - Hober D, *et al.* 2002. *J. Gen. Virol.* 83:2169. (Block)
 - Brainard DM, *et al.* 2009. *J. Virol.* 83:7305. [PubMed](#)
 - Smed-Sørensen A, *et al.* 2008. *Blood* 111:5037. (Block) [PubMed](#)
 - Timmerman KL, *et al.* 2008. *J. Leukoc. Biol.* 84:1271. (FC) [PubMed](#)
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
 - Kim WK, *et al.* 2006. *Am. J. Pathol.* 168:822. (FC)
 - Boltz A, *et al.* 2011. *J. Biol Chem.* 286:21896. [PubMed](#)
 - Wu Z, *et al.* 2013. *J. Virol.* 87:7717. [PubMed](#)
-

Description: CD16 is known as low affinity IgG receptor III (FcγRIII). It is expressed as two distinct forms (CD16a and CD16b). CD16a (FcγRIIIA) is a 50-65 kD polypeptide-anchored transmembrane protein. It is expressed on the surface of NK cells, activated monocytes, macrophages, and placental trophoblasts in humans. CD16b (FcγRIIIB) is a 48 kD glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

- Antigen**
- References:**
- Fleit H, *et al.* 1982. *P. Natl. Acad. Sci. USA* 79:3275.
 - Stroncek D, *et al.* 1991. *Blood* 77:1572.
 - Wirthmueller U, *et al.* 1992. *J. Exp. Med.* 175:1381.