

PE/Dazzle™ 594 anti-human CD16

Catalog # / Size: 2110270 / 100 tests
2110265 / 25 tests

Clone: 3G8

Isotype: Mouse IgG1, κ

Immunogen: Human PMN cells


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V NK80

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD16 (clone 3G8) PE/Dazzle™ 594 (filled histogram) or mouse IgG1, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: The 3G8 antibody blocks neutrophil phagocytosis and stimulates NK cell proliferation. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁶, immunoprecipitation³, stimulation of NK cell proliferation⁴, blocking of phagocytosis⁵, and blocking of immunoglobulin binding to CD16^{7,8}. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 302014). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302050) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

- Application References:**
1. Knapp W, *et al.* Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York.
 2. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 3. Edberg J, *et al.* 1997. *J. Immunol.* 159:3849. (IP)
 4. Hoshino S, *et al.* 1991. *Blood* 78:3232. (Stim)
 5. Tamm A, *et al.* 1996. *Immunol.* 157:1576. (Block)
 6. Da Silva DM, *et al.* 2001. *Int. Immunol.* 13:633. (IHC)
 7. Holl V, *et al.* 2004. *J. Immunol.* 173:6274. (Block)
 8. Hober D, *et al.* 2002. *J. Gen. Virol.* 83:2169. (Block)

9. Brainard DM, *et al.* 2009. *J. Virol.* 83:7305. [PubMed](#)
 10. Smed-Sørensen A, *et al.* 2008. *Blood* 111:5037. (Block) [PubMed](#)
 11. Timmerman KL, *et al.* 2008. *J. Leukoc. Biol.* 84:1271. (FC) [PubMed](#)
 12. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 13. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
 14. Kim WK, *et al.* 2006. *Am. J. Pathol.* 168:822. (FC)
 15. Boltz A, *et al.* 2011. *J. Biol Chem.* 286:21896. [PubMed](#)
 16. Wu Z, *et al.* 2013. *J. Virol.* 87:7717. [PubMed](#)
-

Description: CD16 is known as low affinity IgG receptor III (FcγRIII). It is expressed as two distinct forms (CD16a and CD16b). CD16a (FcγRIIIA) is a 50-65 kD polypeptide-anchored transmembrane protein. It is expressed on the surface of NK cells, activated monocytes, macrophages, and placental trophoblasts in humans. CD16b (FcγRIIIB) is a 48 kD glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

- Antigen**
References:
1. Fleit H, *et al.* 1982. *P. Natl. Acad. Sci. USA* 79:3275.
 2. Stroncek D, *et al.* 1991. *Blood* 77:1572.
 3. Wirthmueller U, *et al.* 1992. *J. Exp. Med.* 175:1381.