

PE/Cyanine5 anti-human CD18

Catalog # / Size: 2110545 / 25 tests

Clone: TS1/18

Isotype: Mouse IgG1, κ


Reactivity: Human, Other

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE/Cy5 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: V AS162

Concentration: Lot-specific




Human peripheral blood lymphocytes stained with TS1/18 PE/Cy5

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of cell adhesion and migration^{3,4}. The LEAF™ Purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 302112). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302116) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).



Formalin-fixed paraffin-embedded human breast cancer tissue slices were deparaffinized and rehydrated. Antigen retrieval was done with Tris-Buffered Saline 1X (1.0 M, pH 7.4) at 95°C for 40 minutes, washed with PBS/0.05% Tween 20 twice for five minutes, permeabilized with 0.5% Triton X-100 for ten minutes, and blocked with 5% FBS and 0.2% gelatin for 30 minutes. Then, the slices were stained with 5 μ g/mL anti-EGFR (clone A19002A) Alexa Fluor® 647 (red) at 4°C overnight. Nuclei were counterstained with DAPI (green). The image was captured with a 10X objective.

**Application
References:**

1. Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
2. Kishimoto T, *et al.* 1997. Leucocyte Typing VI. Garland Press. London.
3. Van Epps DE, *et al.* 1989. *J. Immunol.* 143:3207. (Block)
4. Meerschaert J, *et al.* 1994. *J. Immunol.* 152:1915. (Block)
5. Sithu SD, *et al.* 2007. *J. Biol. Chem.* doi:10.1074/jbc.M611273200.
6. Sommaggio R, *et al.* 2012. *J. Immunol.* 188:2075. [PubMed](#)
7. Valenzuela NM, *et al.* 2013. *J. Immunol.* 190:6635. [PubMed](#)

Description: CD18 is a 90-95 kD type I transmembrane protein also known as integrin β_2 subunit, LFA-1 β subunit, and β_2 integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.

**Antigen
References:**

1. Anderson D, *et al.* 1987. *Annu. Rev. Med.* 38:175.
2. Springer T. 1994. *Cell* 76:301.