

PE/Cy7 anti-human CD18

Catalog # / Size: 2110585 / 25 tests
2110590 / 100 tests

Clone: TS1/18

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V AS162

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD18 (clone TS1/18) PE/Cy7 (filled histogram) or mouse IgG1, κ PE/Cy7 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of cell adhesion and migration^{3,4}. The LEAF™ Purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 302112). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302116) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

- Application References:**
- Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Kishimoto T, *et al.* 1997. Leucocyte Typing VI. Garland Press. London.
 - Van Epps DE, *et al.* 1989. *J. Immunol.* 143:3207. (Block)
 - Meerschaert J, *et al.* 1994. *J. Immunol.* 152:1915. (Block)
 - Sithu SD, *et al.* 2007. *J. Biol. Chem.* doi:10.1074/jbc.M611273200.
 - Sommaggio R, *et al.* 2012. *J. Immunol.* 188:2075. [PubMed](#)
 - Valenzuela NM, *et al.* 2013. *J. Immunol.* 190:6635. [PubMed](#)

Description: CD18 is a 90-95 kD type I transmembrane protein also known as integrin β_2 subunit, LFA-1 β subunit, and β_2 integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.

- Antigen References:**
- Anderson D, *et al.* 1987. *Annu. Rev. Med.* 38:175.
 - Springer T. 1994. *Cell* 76:301.