## **Product Data Sheet**

## APC/Cyanine7 anti-human CD18

**Catalog** # / 2110665 / 25 tests

**Size:** 2110670 / 100 tests

Clone: TS1/18

**Isotype:** Mouse IgG1,  $\kappa$  **Reactivity:** Human, Other

**Preparation:** The antibody was purified by affinity

chromatography and conjugated with

APC/Cyanine7 under optimal

conditions.


**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA)

Workshop Number: V AS162

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with anti-human CD18 (clone TS1/18) APC/Cyanine7 (filled histogram) or mouse IgG1, κ APC/Cyanine7 isotype control (open histogram).

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5  $\mu L$  per million cells in 100  $\mu L$  staining volume or 5  $\mu L$  per 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes:

Additional reported applications (for the relevant formats) include:

inhibition of cell adhesion and migration<sup>3,4</sup>.

Application References:

1. Schlossman S, et al. 1995. Leucocyte Typing V. Oxford University Press. New York.

2. Kishimoto T, et al. 1997. Leucocyte Typing VI. Garland Press. London.

Van Epps DE, et al. 1989. J. Immunol. 143:3207. (Block)
Meerschaert J, et al. 1994. J. Immunol. 152:1915. (Block)

5. Sithu SD, et al. 2007. J. Biol. Chem. doi:10.1074/jbc.M611273200.

6. Sommaggio R, et al. 2012. J. Immunol. 188:2075. PubMed

7. Valenzuela NM, et al. 2013. J. Immunol. 190:6635. PubMed

**Description:** CD18 is a 90-95 kD type I transmembrane protein also known as integrin  $\beta_2$ 

subunit, LFA-1  $\beta$  subunit, and  $\beta_2$  integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18

and associated  $\alpha$  chains function in adhesion and signaling in

hematopoietic cells.

Antigen

1. Anderson D, et al. 1987. Annu. Rev. Med. 38:175.

References:

2. Springer T. 1994. Cell 76:301.