

PE anti-human CD19

Catalog # / Size: 2111040 / 100 tests
2111035 / 25 tests

2111270 / 100 µg

Clone: HIB19

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: microg size: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
test size: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD19.11

Concentration: microg sizes: 0.2 mg/ml
test sizes: lot-specific



Human peripheral blood lymphocytes stained with HIB19 PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining using the microg size, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁸ and blocking of B cell proliferation. Clone HIB19 is not recommended for formalin-fixed paraffin-embedded sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 302214).

- Application References:**
- Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
 - Bradbury L, *et al.* 1993. *J. Immunol.* 151:2915.
 - Joseph A, *et al.* 2010. *J. Virol.* 84:6645. [PubMed](#)
 - Wang X, *et al.* 2010. *Haematologica.* 95:884. (FC) [PubMed](#)
 - Walker JD, *et al.* 2009. *J. Immunol.* 182:1548. (Block) [PubMed](#)
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Hansen A, *et al.* 2002. *Arthritis Rheum.* 46:2160. (IHC)
 - White CA, *et al.* 2014. *J Immunol.* 193:5933. [PubMed](#)

Description: CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cells, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.

Antigen 1. Tedder T, *et al.* 1994. *Immunol. Today* 15:437.
References: 2. Bradbury L, *et al.* 1993. *J. Immunol.* 151:2915.