

PerCP/Cy5.5 anti-human CD19

Catalog # / Size: 2111145 / 25 tests
2111150 / 100 tests

Clone: HIB19

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD19.11

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with HIB19 PerCP/Cy5.5

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells in 100 microL volume or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁸ and blocking of B cell proliferation. Clone HIB19 is not recommended for formalin-fixed paraffin-embedded sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 302214).

Application References:

- Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
- Bradbury L, *et al.* 1993. *J. Immunol.* 151:2915.
- Joseph A, *et al.* 2010. *J. Virol.* 84:6645. [PubMed](#)
- Wang X, *et al.* 2010. *Haematologica.* 95:884. (FC) [PubMed](#)
- Walker JD, *et al.* 2009. *J. Immunol.* 182:1548. (Block) [PubMed](#)
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Hansen A, *et al.* 2002. *Arthritis Rheum.* 46:2160. (IHC)
- Lepore M, *et al.* 2014. *J Exp Med.* 211:1363. [PubMed](#)
- Garcia-Carmona Y, *et al.* 2015. *Blood.* 125:1749. [PubMed](#)
- Weinberg A, *et al.* 2015. *PLoS One.* 10:122431. [PubMed](#)

Description: CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a

member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cells, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.

- Antigen**
- References:**
1. Tedder T, *et al.* 1994. *Immunol. Today* 15:437.
 2. Bradbury L, *et al.* 1993. *J. Immunol.* 151:2915.