

APC/Fire™ 750 anti-human CD20

Catalog # / Size: 2111790 / 100 tests
2111785 / 25 tests

Clone: 2H7

Isotype: Mouse IgG2b, κ

Immunogen: Human tonsillar B cells


Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: 750 under optimal conditions.

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 PE and CD20 (clone 2H7) APC/Fire™ 750 (top) or mouse IgG2b, κ APC/Fire™ 750 isotype control (bottom).


Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: The epitope recognized by clone 2H7 has been mapped to the sequence YNCEPANPSEKNSPST which lies in the large extracellular loop of human CD20. Additional reported applications (for the relevant formats) include:
immunoprecipitation⁴ and immunohistochemical staining of acetone-fixed frozen sections⁵.

- Application References:**
- Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
 - McMichael A, *et al.* Eds. 1987. Leucocyte Typing III Oxford University Press. New York.
 - Polyak MJ, *et al.* 2002. *Blood* 99:3256. (IP)
 - Mack CL, *et al.* 2004. *Pediatr. Res.* 56:79. (IHC)



Description: CD20 is a 33-37 kD, four transmembrane spanning protein, also known as B1 and Bp35. CD20 is expressed on pre-B-cells, resting and activated B cells (not plasma cells), some follicular dendritic cells, and at low levels on a T cell subset. CD20 is heavily phosphorylated on activated B cells and malignant B cells. Homo-oligomeric complexes of CD20 are thought to form Ca^{2+} conductive ion channels in the plasma membrane of B cells. The CD20 molecule is involved in B-cell activation and is associated with various Src family kinases (Lyn, Lck, Fyn). It exists in a complex with MHC class I and II, CD53, CD81, and CD82.

Antigen 1. Hultin L, *et al.* 1993. *Cytometry* 14:196.
References: 2. Tedder T, *et al.* 1994. *Immunol. Today* 15:450.