


APC anti-human CD22

Catalog # / Size: 2112550 / 100 tests
Clone: HIB22
Isotype: Mouse IgG1, κ
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Workshop Number: V CD22.14
Concentration: NULL



Human peripheral blood lymphocytes stained with HIB22 APC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections.

- Application References:**
- Schlossman S, et al. Eds. 1995. Leukocyte Typing V:White Cell Differentiation Antigens. Oxford University Press. New York.
 - Clark E. 1993. *J. Immunol.* 150:4715.
 - Shan D and O. Press. 1995. *J. Immunol.* 154:4466.
 - Herter S, et al. 2013. *Mol Cancer ther.* 12:2031. [PubMed](#)
 - Rossi EA, et al. 2014. *PLoS One.* 9:98315. [PubMed](#)

Description: CD22 is a 130 kD type I transmembrane glycoprotein also known as Siglec-2 and BL-CAM. It is a member of the immunoglobulin superfamily (sialoadhesion subgroup). CD22 is expressed in the cytoplasm of pro-B and pre-B cells, and on the surface of mature B and activated B cells, but not on plasma cells. CD22 is present in the B cell receptor complex and associates with SHP-1, Syk, Lck, Lyn, and phospholipase Cy1. A primary function of CD22 is thought to be in limiting antigen receptor signaling by modulating B cell activation threshold. CD22 has been shown to bind to CD45RO and CD75, although the natural ligands for this molecule remain controversial.

- Antigen References:**
- Clark E. 1993. *J. Immunol.* 150:4715.
 - Shan D, et al. 1995. *J. Immunol.* 154:4466.