


FITC anti-human CD25

Catalog # / Size:	2113020 / 100 tests 2113015 / 25 tests
Clone:	BC96
Isotype:	Mouse IgG1, κ
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Workshop Number:	V T-072
Concentration:	Lot-specific



PHA-stimulated (3 day) human peripheral blood lymphocytes were stained with CD25 (clone BC96) FITC (filled histogram) or mouse IgG1, κ FITC isotype control (open histogram).

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 microL to 5 microL per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	Additional reported applications include: immunofluorescence ³ .
Application References:	<ol style="list-style-type: none">Schlossman S, <i>et al.</i> Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.Ernst CW, <i>et al.</i> 2007. <i>Clin. Exp. Immunol.</i> 148:271. (IF) PubMedKmieciak M, <i>et al.</i> 2009. <i>J. Transl. Med.</i> 7:89. (FC) PubMedBusinaro R, <i>et al.</i> 2009. <i>Altherosclerosis.</i> 207:74. PubMedRyba M, <i>et al.</i> 2011. <i>Cytokine.</i> 55:353. PubMed

Description: CD25 is a 55 kD type I transmembrane glycoprotein also known as the low affinity IL-2 receptor α chain or Tac. It is expressed on progenitor lymphocytes, activated T and B cells, and activated monocytes/macrophages. CD25 is also expressed on a subset of non-stimulated CD4⁺ T cells termed T regulatory cells. CD25 associates with the IL-2 receptor β (CD122) and common γ chains (CD132) to form the high affinity IL-2R complex.

Antigen References:

- Taniguchi T, *et al.* 1993. *Cell* 73:5.
- Waldmann T. 1991. *J. Biol. Chem.* 266:2681.