
Product Data Sheet

APC anti-human CD25

Catalog # / Size:	2113050 / 100 tests 2113045 / 25 tests	
Clone:	BC96	
Isotype:	Mouse lgG1, κ	all Number
Reactivity:	Human	
Preparation:	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.	Relative Co
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	10 ⁰ PHA-sti
Workshop Number:	V T-072	periph stained
Concentration:	Lot-specific	BC96) / mouse



PHA-stimulated (3 day) human peripheral blood lymphocytes were stained were with CD25 (clone BC96) APC (filled histogram) or mouse lgG1, κ APC isotype control (open histogram).

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 microL to 5 microL per test . Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	Additional reported applications include: immunofluorescence3.
Application References:	 Schlossman S, <i>et al.</i> Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. Kmieciak M, <i>et al.</i> 2009. <i>J. Transl. Med.</i> 7:89. (FC) <u>PubMed</u> Ernst CW, <i>et al.</i> 2007. <i>Clin. Exp. Immunol.</i> 148:271. (IF) <u>PubMed</u> Trabanelli S, <i>et al.</i> 2014. <i>J Immunol.</i> 192:1231. <u>PubMed</u>
Description:	CD25 is a 55 kD type I transmembrane glycoprotein also known as the low affinity IL-2 receptor α chain or Tac. It is expressed on progenitor lymphocytes, activated T and B cells, and activated monocytes/macrophages. CD25 is also expressed on a subset of non-stimulated CD4 ⁺ T cells termed T regulatory cells. CD25 associates with the IL-2 receptor β (CD122) and common γ chains (CD132) to form the high affinity IL-2R complex.
Antigen References:	1. Taniguchi T, <i>et al.</i> 1993. <i>Cell</i> 73:5. 2. Waldmann T. 1991. <i>J. Biol. Chem.</i> 266:2681.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com