

Alexa Fluor® 647 anti-human CD29

Catalog # / Size: 2115085 / 25 tests
2115090 / 100 tests

Clone: TS2/16

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V A-S202

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with TS2/16 Alexa Fluor® 647

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation³, immunohistochemical staining of acetone-fixed frozen tissue sections^{3,5}, and activation of integrin β₁^{4,7,8}. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 303010). Clone TS2/16 recognizes epitope A2.¹⁰

- Application References:**
1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 2. Gutierrez-Lopez M, *et al.* 2003. *J. Biol. Chem.* 278:208.
 3. Hemler ME, *et al.* 1984. *J. Immunol.* 132:3011. (IHC, IP)
 4. Sanchez-Aparicio P, *et al.* 1994. *J. Cell Biol.* 126:271. (Activ)
 5. Frank NY, *et al.* 2005. *Cancer Res.* 65:4320. (IHC)
 6. Murga M, *et al.* 2005. *Blood* 105:1992. (FC) [PubMed](#)
 7. Porter JC and Hogg N. 1997. *J. Cell Biol.* 138:1437. (Activ)
 8. Conway RE, *et al.* 2006. *Mol. Cell. Biol.* 26:5310. (Activ)
 9. Wesseling J, *et al.* 1995. *J. Cell. Biol.* 129:255. (Dog Reactivity)
 10. Rubio G, *et al.* 2002. *Cancer Immunol. Immunother.* 51:130.
 11. Dong A, *et al.* 2015. *J Biol Chem.* 290:8016. PubMed
 12. Paebst F, *et al.* 2014. *Cytometry A.* 85(8):678-87. (Horse reactivity)

Description: CD29 is a 130 kD single chain type I glycoprotein also known as integrin β₁, VLA-β chain, or gp11a. It is broadly expressed on a majority of hematopoietic and non-

hematopoietic cells, including leukocytes (although at low level on granulocytes), platelets, fibroblasts, endothelial cells, epithelial cells, and mast cells. CD29 is a member of the integrin family. It is non-covalently associated with integrin α 1- α 6 chains to form VLA-1 to VLA-6 molecules, respectively. Integrins, which include CD29, bind to several cell surface (e.g. VCAM-1, MadCAM-1) and extracellular matrix molecules. CD29 acts as a fibronectin receptor and is involved in a variety of cell-cell and cell-matrix interactions.

- Antigen**
References:
1. Hemler M. 1990. *Annu. Rev. Immunol.* 8:365.
 2. Hynes R. 1992. *Cell* 69:11.