
Product Data Sheet

Purified anti-human CD38

Catalog # / Size:	2117510 / 100 µg
Clone:	HIT2
Isotype:	Mouse IgG1, κ
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Workshop Number:	III 155
Concentration:	0.5



Human peripheral blood lymphocytes stained with purified HIT2, followed by anti-mouse IgGs FITC

Applications:

Applications:	Flow Cytometry, Immunohistochemistry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is \leq 2.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application	Additional reported applications (for the relevant formats) include:
Notes:	immunohistochemical staining of acetone-fixed frozen tissue sections. ⁶
Application	 Kishimoto T, <i>et al.</i> Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc.
References:	London. Dieu M. 1998. <i>J. Exp. Med.</i> 188:373. Esser M, <i>et al.</i> 2001. <i>J. Virol.</i> 75:6173. Jeannin P, <i>et al.</i> 1999. <i>J. Immunol.</i> 162:2044. Kapsogeorgou EK, <i>et al.</i> 2001. <i>J. Immunol.</i> 166:3107. van der Voort R, <i>et al.</i> 1997. <i>J. Exp. Med.</i> 185:2121. (IHC) Bende RJ, <i>et al.</i> 2003. <i>Am. J. Pathol.</i> 162:105. Lehner M, <i>et al.</i> 2008. <i>J. Leukoc. Biol.</i> 83:883. PubMed Yoshino N, <i>et al.</i> 2000. <i>Exp. Anim. (Tokyo)</i> 49:97. (FC)
Description:	CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.
Antigen	1. Ferrero E, <i>et al.</i> 1999. <i>J. Leukoc. Biol.</i> 65:151.
References:	2. Lund F, <i>et al.</i> 1995. <i>Immunol. Today</i> 16:469.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com