

PerCP/Cy5.5 anti-human CD38

Catalog # / Size: 2117605 / 25 tests
2117610 / 100 tests

Clone: HIT2

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: III 155

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with HIT2 PerCP/Cy5.5

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections.⁶

- Application References:**
1. Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
 2. Dieu M. 1998. *J. Exp. Med.* 188:373.
 3. Esser M, *et al.* 2001. *J. Virol.* 75:6173.
 4. Jeannin P, *et al.* 1999. *J. Immunol.* 162:2044.
 5. Kapsogeorgou EK, *et al.* 2001. *J. Immunol.* 166:3107.
 6. van der Voort R, *et al.* 1997. *J. Exp. Med.* 185:2121. (IHC)
 7. Bende RJ, *et al.* 2003. *Am. J. Pathol.* 162:105.
 8. Lehner M, *et al.* 2008. *J. Leukoc. Biol.* 83:883. [PubMed](#)
 9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)

Description: CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

- Antigen** 1. Ferrero E, *et al.* 1999. *J. Leukoc. Biol.* 65:151.
References: 2. Lund F, *et al.* 1995. *Immunol. Today* 16:469.