
Product Data Sheet

Brilliant Violet 605™ anti-human CD41

Catalog # / Size:	2118710 / 100 tests 2118705 / 25 tests	□ Human peripheral blood platelets were stained with anti-human CD41 (clone HIP8) Brilliant Violet 605™ (filled histogram) or mouse IgG1, κ Brilliant Violet 605™ isotype control (open histogram)
Clone:	HIP8	
Isotype:	Mouse IgG1, κ	
Immunogen:	Mouse IL-9R transfectants	
Reactivity:	Human, Non-human primate, Other	
Preparation:	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 605™ and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)	
Workshop Number:	IV P38	
Concentration:	Lot-specific	

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections and blocking of platelet aggregation². The HIP8 antibody has been reported to block the activation of platelets by various stimuli, including collagen, and ADP.

**Application
References:**

1. Knapp W, et al. 1989. Leucocyte Typing IV. Oxford University Press. New York.
 2. McCarty OJT, et al. 2000. *Blood* 96:1789.
 3. Yoshino N, et al. 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 4. Zhi L et al. 2013. *PLoS One*. 8:e79869. (IHC)
-

Description: CD41 is a 125/25 kD α subunit of the gpIIb/IIIa (CD41/CD61) complex. CD41 is a heterodimer composed of a heavy chain (gpIIb α) and light chain (gpIIb β) linked by a single disulfide bond. It is a member of the integrin family primarily expressed on platelets and megakaryocytes. CD41 has been reported to be involved with platelet aggregation and platelet attachment to the ECM. CD41/CD61 complex acts as the receptor for fibrinogen, fibronectin, Von Willebrand factor, and thrombin.

**Antigen
References:**

1. Denzin L, et al. 1996. *J. Exp. Med.* 184:2153.
2. Denzin L, et al. 1995. *Cell* 82:155.
3. Riberdy J, et al. 1994. *J. Cell Biol.* 125:1225.