


FITC anti-human CD42b

Catalog # / Size: 2119515 / 25 tests
Clone: HIP1
Isotype: Mouse IgG1, κ
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Workshop Number: IV P70
Concentration: Lot-specific



Human peripheral blood platelets stained with purified HIP1 FITC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone HIP1 recognizes an epitope within the N-terminal region of the GPIIb/IIIa chain. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections, Western blotting, and inhibition of platelet aggregation. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 303908).

- Application References:**
- Knapp W, *et al.* 1989. *Leucocyte Typing IV*. Oxford University Press. New York.
 - Takahashi R, *et al.* 1999. *Blood*. 93:1951.
 - Saggu G, *et al.* 2013. *J. Immunol.* 190:6457. [PubMed](#)
 - Meyer Dos Santos S, *et al.* 2011. *Blood*. 117:4999. (Block) [PubMed](#)
 - Vettore S, *et al.* 2008. *Haematologica*. 93:1743.

Description: CD42b is a 145 kD glycoprotein known as gplIb. It is covalently bonded to CD42c to form GPIIb. CD42b is expressed on platelets and megakaryocytes. CD42b/c heterodimer forms a complex with CD42a and d and acts as the receptor for von Willibrand factor and thrombin.

- Antigen References:**
- Clemetson K, *et al.* 1982. *J. Clin. Invest.* 70:304.
 - Fox J, *et al.* 1988. *J. Biol. Chem.* 263:4882.
 - Kuijpers R, *et al.* 1992. *Blood* 79:283.