


PerCP/Cy5.5 anti-human CD42b

Catalog # / Size:	2119590 / 100 tests
Clone:	H1P1
Isotype:	Mouse IgG1, κ
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Workshop Number:	IV P70
Concentration:	0.5



Human platelets were stained with CD42b (clone H1P1) PerCP/Cy5.5 (filled histogram) or mouse IgG1, κ PerCP/Cy5.5 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Clone H1P1 recognizes an epitope within the N-terminal region of the GPIIb/IIIa chain. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections, Western blotting, and inhibition of platelet aggregation. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 303908).

Application References:

- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
- Takahashi R, *et al.* 1999. *Blood*. 93:1951.
- Saggu G, *et al.* 2013. *J. Immunol.* 190:6457. [PubMed](#)
- Meyer Dos Santos S, *et al.* 2011. *Blood*. 117:4999. (Block) [PubMed](#)
- Vettore S, *et al.* 2008. *Haematologica*. 93:1743.

Description: CD42b is a 145 kD glycoprotein known as GPIIb/IIIa. It is covalently bonded to CD42c to form GPIIb/IIIa. CD42b is expressed on platelets and megakaryocytes. CD42b/c heterodimer forms a complex with CD42a and d and acts as the receptor for von Willibrand factor and thrombin.

Antigen References:

- Clemetson K, *et al.* 1982. *J. Clin. Invest.* 70:304.
- Fox J, *et al.* 1988. *J. Biol. Chem.* 263:4882.
- Kuijpers R, *et al.* 1992 *Blood* 79:283.