

Brilliant Violet 421™ anti-human CD42b

Catalog # / Size: 2119645 / 25 tests
2119650 / 100 tests

Clone: HIP1

Isotype: Mouse IgG1, κ

Reactivity: Human

Concentration: Lot-specific

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Clone HIP1 recognizes an epitope within the N-terminal region of the GPIIb α chain5. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections, Western blotting, and inhibition of platelet aggregation2. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 303908).

Description: CD42b is a 145 kD glycoprotein known as gplb α . It is covalently bonded to CD42c to form GPIIb. CD42b is expressed on platelets and megakaryocytes. CD42b/c heterodimer forms a complex with CD42a and d and acts as the receptor for von Willibrand factor and thrombin.