


Biotin anti-human CD45RA

Catalog # / Size: 2120520 / 100 µg
Clone: HI100
Isotype: Mouse IgG2b, κ
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Workshop Number: IV N906
Concentration: 0.5



Human peripheral blood lymphocytes stained with biotinylated HI100, followed by Sav-PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 microg per 10^6 cells in 100 microL volume or 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for relevant formats of this clone) include: inhibition of CD45 functions², immunohistochemical staining of frozen tissue sections³ and formalin-fixed paraffin-embedded tissue sections⁴, and immunofluorescence^{15,16}.

- Application References:**
- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
 - Yamada T, *et al.* 2002. *J. Biol. Chem.* 277:28830. (WB, Block)
 - Weninger W, *et al.* 2003 *J. Immunol.* 170:4638. (IHC)
 - Imanguli MM, *et al.* 2009. *Blood.* 113:3620 (IHC)
 - Roque S, *et al.* 2007. *J. Immunol.* 178:8028. (FC) [PubMed](#)
 - Smeltz RB. 2007. *J. Immunol.* 178:4786. (FC) [PubMed](#)
 - Palendira U, *et al.* 2008. *Blood* (FC) [PubMed](#)
 - Kuttruff S, *et al.* 2009. *Blood* 113:358. (FC) [PubMed](#)
 - Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
 - Alanio C, *et al.* 2010. *Blood* 115:3718. (FC) [PubMed](#)
 - Iannello A, *et al.* 2010. *J. Immunol.* 184:114. (FC) [PubMed](#)
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Guereau-de-Arellan M, *et al.* 2011. *Brain.* 134:3578. [PubMed](#)
 - Canque B, *et al.* 2000. *Blood* 96:3748. (IF)
 - Imanguli MM, *et al.* 2009. *Blood* 13:3620. (IF)

Description: CD45RA is a 205-220 kD single chain type I glycoprotein. It is an exon 4 splice variant of the tyrosine phosphatase CD45. The CD45RA isoform is expressed on resting/naïve T cells, medullary thymocytes, B cells and monocytes. CD45RA enhances both T cell receptor and B cell receptor signaling. CD45 non-covalently

associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1. CD45 isoform expression can change in response to cytokines.

Antigen
References:

1. Thomas M. 1989. *Annu. Rev. Immunol.* 7:339.
2. Trowbridge I, *et al.* 1994. *Annu. Rev. Immunol.* 12:85.