

# Alexa Fluor® 700 anti-human CD45RA

**Catalog # /** 2120600 / 100 µg  
**Size:** 2120595 / 25 µg

**Clone:** HI100

**Isotype:** Mouse IgG2b, κ


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 700 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Workshop Number:** IV N906

**Concentration:** 0.5



Human peripheral blood  
 lymphocytes stained with HI100  
 Alexa Fluor® 700

## Applications:

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is highly recommended that the reagent be titrated for optimal performance for each application.

\* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

**Application Notes:** Additional reported applications (for relevant formats of this clone) include: inhibition of CD45 functions<sup>2</sup>, immunohistochemical staining of frozen tissue sections<sup>3</sup> and formalin-fixed paraffin-embedded tissue sections<sup>4</sup>, and immunofluorescence<sup>15,16</sup>.

- Application References:**
1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
  2. Yamada T, *et al.* 2002. *J. Biol. Chem.* 277:28830. (WB, Block)
  3. Weninger W, *et al.* 2003 *J. Immunol.* 170:4638. (IHC)
  4. Imanguli MM, *et al.* 2009. *Blood.* 113:3620 (IHC)
  5. Roque S, *et al.* 2007. *J. Immunol.* 178:8028. (FC) [PubMed](#)
  6. Smeltz RB. 2007. *J. Immunol.* 178:4786. (FC) [PubMed](#)
  7. Palendira U, *et al.* 2008. *Blood* (FC) [PubMed](#)
  8. Kutruff S, *et al.* 2009. *Blood* 113:358. (FC) [PubMed](#)
  10. Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
  11. Alanio C, *et al.* 2010. *Blood* 115:3718. (FC) [PubMed](#)
  12. Iannello A, *et al.* 2010. *J. Immunol.* 184:114. (FC) [PubMed](#)
  13. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
  14. Guereau-de-Arellan M, *et al.* 2011. *Brain.* 134:3578. [PubMed](#)
  15. Canque B, *et al.* 2000. *Blood* 96:3748. (IF)
  16. Imanguli MM, *et al.* 2009. *Blood* 113:3620. (IF)

- Description:** CD45RA is a 205-220 kD single chain type I glycoprotein. It is an exon 4 splice variant of the tyrosine phosphatase CD45. The CD45RA isoform is expressed on resting/naïve T cells, medullary thymocytes, B cells and monocytes. CD45RA enhances both T cell receptor and B cell receptor signaling. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1. CD45 isoform expression can change in response to cytokines.
- Antigen** 1. Thomas M. 1989. *Annu. Rev. Immunol.* 7:339.
- References:** 2. Trowbridge I, *et al.* 1994. *Annu. Rev. Immunol.* 12:85.