

Brilliant Violet 650™ anti-human CD45RA

Catalog # / Size: 2120675 / 25 tests
2120680 / 100 tests

Clone: HI100

Isotype: Mouse IgG2b, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 650™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 650™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: IV N906

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD45RO FITC and CD45RA (clone HI100) Brilliant Violet 650™.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 650™ excites at 405 nm and emits at 645 nm. The bandpass filter 660/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 650™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for relevant formats of this clone) include: inhibition of CD45 functions², immunohistochemical staining of frozen tissue sections³ and formalin-fixed paraffin-embedded tissue sections⁴, and immunofluorescence^{15,16}.

Application References:

1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
2. Yamada T, *et al.* 2002. *J. Biol. Chem.* 277:28830. (WB, Block)
3. Weninger W, *et al.* 2003 *J. Immunol.* 170:4638. (IHC)
4. Imanguli MM, *et al.* 2009. *Blood.* 113:3620 (IHC)
5. Roque S, *et al.* 2007. *J. Immunol.* 178:8028. (FC) [PubMed](#)

6. Smeltz RB. 2007. *J. Immunol.* 178:4786. (FC) [PubMed](#)
 7. Palendira U, *et al.* 2008. *Blood* (FC) [PubMed](#)
 8. Kuttruff S, *et al.* 2009. *Blood* 113:358. (FC) [PubMed](#)
 10. Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
 11. Alanio C, *et al.* 2010. *Blood* 115:3718. (FC) [PubMed](#)
 12. Iannello A, *et al.* 2010. *J. Immunol.* 184:114. (FC) [PubMed](#)
 13. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 14. Guereau-de-Arellan M, *et al.* 2011. *Brain.* 134:3578. [PubMed](#)
 15. Canque B, *et al.* 2000. *Blood* 96:3748. (IF)
 16. Imanguli MM, *et al.* 2009. *Blood* 13:3620. (IF)
-

Description: CD45RA is a 205-220 kD single chain type I glycoprotein. It is an exon 4 splice variant of the tyrosine phosphatase CD45. The CD45RA isoform is expressed on resting/naïve T cells, medullary thymocytes, B cells and monocytes. CD45RA enhances both T cell receptor and B cell receptor signaling. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1. CD45 isoform expression can change in response to cytokines.

Antigen 1. Thomas M. 1989. *Annu. Rev. Immunol.* 7:339.
References: 2. Trowbridge I, *et al.* 1994. *Annu. Rev. Immunol.* 12:85.