


FITC anti-human CD49d

Catalog # / Size:	2121580 / 100 tests 2121575 / 25 tests
Clone:	9F10
Isotype:	Mouse IgG1, κ
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Workshop Number:	V S215
Concentration:	Lot-specific



Human peripheral blood lymphocytes were stained with CD49d (clone 9F10) FITC (filled histogram) or mouse IgG1, κ FITC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections, and *in vitro* T cell costimulation^{2,3}. The LEAF™ Purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 304310).

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Jeong SH, *et al.* 2004. *J. Virol.* 78:6995. (Costim)
 - Vogel TU, *et al.* 2002. *J. Immunol.* 169:4511. (Costim)
 - Kleinewietfeld M, *et al.* 2009. *Blood* 113:827. (FC) [PubMed](#)
 - Palacios F, *et al.* 2010. *Blood* 115:4488. [PubMed](#)
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Sestak K, *et al.* 2007. *Vet. Immunol. Immunopathol.* 119:21.
 - Mattapallil MJ, *et al.* 2011. *J. Immunol.* 187:1977. [PubMed](#)

Description: CD49d is a 150 kD α integrin chain known as α_4 integrin or VLA-4 α chain. It forms a heterodimer with either integrin β_1 ($\alpha_4\beta_1$, VLA-4) or β_7 ($\alpha_4\beta_7$). CD49d is expressed broadly on T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils, mast cells, NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red blood cells, platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin. $\alpha_4\beta_7$ is the receptor for VCAM-1 and MAdCAM-1. CD49d participates in mononuclear cell trafficking to endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to

extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3⁺ cells.

Antigen
References:

1. Elices M, Ed.1995. *Springer Semin. Immunopathol.* 16(4).
2. Lobb RR and Helmer ME. *et al.* 1994. *J. Clin. Invest.* 94:1722.