

PE/Dazzle™ 594 anti-human CD62L

Catalog # / Size: 2124205 / 25 tests
2124210 / 100 tests

Clone: DREG-56

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V S056

Concentration: 0.5



Human peripheral blood lymphocytes were stained with CD62L (clone DREG-56) PE/Dazzle™ 594 (filled histogram) or mouse IgG1, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting^{2,3,9} and *in vitro* blocking of lymphocytes binding to high endothelial venules (HEV)². The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 304812).

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Kishimoto TK, *et al.* 1990. *Proc. Natl. Acad. Sci. USA* 87:2244. (WB, Block)
 - Jutila M, *et al.* 2002. *J. Immunol.* 169:1768. (WB)
 - Tamassia N, *et al.* 2008. *J. Immunol.* 181:6563. (FC) [PubMed](#)
 - Kmieciak M, *et al.* 2009. *J. Transl. Med.* 7:89. (FC) [PubMed](#)
 - Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
 - Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Koenig JM, *et al.* 1996. *Pediatr. Res.* 39:616. (WB)
 - Shi C, *et al.* 2011. *J. Immunol.* 187:5293. (FC) [PubMed](#)
 - Burges M, *et al.* 2013. *Clin Cancer Res.* 19:5675. [PubMed](#)
 - Cash JL, *et al.* 2013. *EMBO Rep.* 14:999. (FC) [PubMed](#)

Description: CD62L is a 74-95 kD single chain type I glycoprotein referred to as L-selectin or LECAM-1. It is expressed on most peripheral blood B cells, subsets of T and NK

cells, monocytes, granulocytes, and certain hematopoietic malignant cells. CD62L binds to carbohydrates present on certain glycoforms of CD34, glycam-1, and MAdCAM-1 and with a low affinity to anionic oligosaccharide sequences related to sialylated Lewis X (sLex, CD15s) through its C-type lectin domain. CD62L is important for the homing of naïve lymphocytes to high endothelial venules in peripheral lymph nodes and Peyer's patches. It also plays a role in leukocyte rolling on activated endothelial cells.

- Antigen**
- References:**
1. Kishimoto T, *et al.* 1990. *P. Natl. Acad. Sci. USA* 87:2244.
 2. Kishimoto T, *et al.* 1991. *Blood* 78:805.