

FITC anti-human CD62P (P-Selectin)

Catalog # / Size: 2124515 / 25 tests
2124520 / 100 tests

Clone: AK4

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: VI P-44

Concentration: Lot-specific



Thrombin-activated human peripheral blood platelets stained with AK4 FITC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴ and *in vitro* blocking of adhesion of platelets¹. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 304911).

- Application References:**
1. Skinner M, *et al.* 1991. *J. Biol. Chem.* 266:5371. (Block)
 2. Kishimoto T, *et al.* Eds. 1997. *Leucocyte Typing VI*. Garland Publishing Inc. London.
 3. Yen YT, *et al.* 2006. *J. Virol.* 80:2684.
 4. Sato Y, *et al.* 2005. *Blood* 106:428. (IHC)

Description: CD62P is a 140 kD type I transmembrane glycoprotein also known as P-selectin, platelet activation-dependent granule membrane protein (PADGEM), and GMP-140. It is expressed on activated platelets, megakaryocytes, and endothelial cells. CD62P is primarily stored in secretory α -granules in platelets and Weibel-Palade bodies in endothelial cells, and is rapidly relocated to the plasma membrane upon activation. The ligands for CD62P are CD162 and CD24. A primary function of CD62P is cell adhesion during neutrophil rolling, and platelet-neutrophil and platelet-monocyte interactions.

- Antigen References:**
1. McEver R, *et al.* 1995. *J. Biol. Chem.* 270:11025.
 2. Varki A. 1994. *P. Natl. Acad. Sci. USA* 91:7390.