

Brilliant Violet 650™ anti-human CD62P (P-Selectin)

Catalog # / Size: 2124690 / 100 tests
2124685 / 25 tests

Clone: AK4

Isotype: Mouse IgG1, κ

Immunogen: Recombinant mouse CD163 extracellular domain


Reactivity: Human, Non-human primate

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 650™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 650™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: VI P-44

Concentration: Lot-specific



Thrombin-activated human peripheral blood platelets were stained with CD62P (P-Selectin) (clone AK4) Brilliant Violet 650™ (filled histogram) or mouse IgG1, κ Brilliant Violet 650™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

Brilliant Violet 650™ excites at 405 nm and emits at 645 nm. The bandpass filter 660/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 650™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴ and *in vitro* blocking of adhesion of platelets¹.

Application**References:**

1. Skinner M, *et al.* 1991. *J. Biol. Chem.* 266:5371. (Block)
 2. Kishimoto T, *et al.* Eds. 1997. *Leucocyte Typing VI*. Garland Publishing Inc. London.
 3. Yen YT, *et al.* 2006. *J. Virol.* 80:2684.
 4. Sato Y, *et al.* 2005. *Blood* 106:428. (IHC)
-

Description:

CD62P is a 140 kD type I transmembrane glycoprotein also known as P-selectin, platelet activation-dependent granule membrane protein (PADGEM), and GMP-140. It is expressed on activated platelets, megakaryocytes, and endothelial cells. CD62P is primarily stored in secretory α -granules in platelets and Weibel-Palade bodies in endothelial cells, and is rapidly relocated to the plasma membrane upon activation. The ligands for CD62P are CD162 and CD24. A primary function of CD62P is cell adhesion during neutrophil rolling, and platelet-neutrophil and platelet-monocyte interactions.

**Antigen
References:**

1. McEver R, *et al.* 1995. *J. Biol. Chem.* 270:11025.
2. Varki A. 1994. *P. Natl. Acad. Sci. USA* 91:7390.