Product Data Sheet

PerCP/Cy5.5 anti-human CD66b

Catalog # / Size: 2125540 / 100 tests

2125535 / 25 tests

Clone: G10F5

Isotype: Mouse IgM, κ

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated

antibody.


Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Workshop Number: VI MA81

Concentration: Lot-specific



Human peripheral blood granulocytes stained with G10F5 PerCP/Cy5.5

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes:

Additional reported applications (for the relevant formats) include:

immunohistochemical staining of acetone-fixed frozen and formalin-fixed paraffin-

embedded tissue sections.

Application References:

1. Schlossman S, et al. Eds. 1995. Leucocyte Typing V. Oxford University Press.

New York.

2. Kishimoto T, et al. Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc.

London.

3. Norling LV, et al. 2012. Arterioscler Thromb Vasc Biol. 32:1970. PubMed

4. Meinke P, et al. 2015. Neuroimmunol Discord. 25:127. PubMed

Description: CD66b is a 95-100 kD glycosylphosphatidylinositol (GPI)-linked protein also known

as CD67, CGM6, and NCA-95. CD66b is a member of the immunoglobulin

superfamily, carcinoembryonic antigen (CEA)-like subfamily. CD66b, expressed on granulocytes, has been reported to induce activation in neutrophils and to be

involved in heterophilic adhesion with CD66c.

Antigen

1. Kuijpers T, et al. 1993. J. Immunol. 151:4934.

References: 2. Kuroki M, et al. 1992. J. Leuk. Biol. 52:551.