

Alexa Fluor® 647 anti-human CD66b

Catalog # / Size: 2125550 / 100 tests
2125545 / 25 tests

Clone: G10F5

Isotype: Mouse IgM, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: VI MA81

Concentration: Lot-specific




Human peripheral blood granulocytes were stained with CD66b (clone G10F5) Alexa Fluor® 647 (filled histogram) or mouse IgM, κ Alexa Fluor® 647 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. For immunohistochemical staining on formalin-fixed paraffin-embedded tissue sections, the suggested use of this reagent is 5.0 - 10 microg per ml. It is recommended that the reagent be titrated for optimal performance for each application.



Human paraffin-embedded spleen tissue was stained with Alexa Fluor® 647 anti-human CD66b antibody (red) and co-stained with Alexa Fluor® 594 anti-human CD8a antibody (green). See additional supplemental data for detailed information.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen and formalin-fixed paraffin-embedded tissue sections.

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
 - Norling LV, *et al.* 2012. *Arterioscler Thromb Vasc Biol.* 32:1970. [PubMed](#)
 - Meinke P, *et al.* 2015. *Neuroimmunol Discord.* 25:127. [PubMed](#)

Description: CD66b is a 95-100 kD glycosylphosphatidylinositol (GPI)-linked protein also known as CD67, CGM6, and NCA-95. CD66b is a member of the immunoglobulin superfamily, carcinoembryonic antigen (CEA)-like subfamily. CD66b, expressed on granulocytes, has been reported to induce activation in neutrophils and to be involved in heterophilic adhesion with CD66c.

Antigen 1. Kuijpers T, *et al.* 1993. *J. Immunol.* 151:4934.
References: 2. Kuroki M, *et al.* 1992. *J. Leuk. Biol.* 52:551.