

PE anti-human CD80

Catalog # / Size: 2126035 / 25 tests
2126040 / 100 tests

Clone: 2D10

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: VI CD80.1

Concentration: Lot-specific



Human B-cell Burkitt's lymphoma cell line Daudi stained with 2D10 PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: *in vitro* blocking of T cell activation, immunohistochemical staining of acetone-fixed frozen tissue sections², immunoprecipitation, and Western blotting³. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 305212).

- Application References:**
1. Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
 2. Battifora M. 1998. *J. Clin. Endocr. Metab.* 83:4130. (IHC)
 3. Van der Merwe PA, *et al.* 1997. *J. Exp. Med.* 185:3. (WB)
 4. Jayakumar A, *et al.* 2008. *Infect. Immun.* 76:2138. [PubMed](#)
 5. Schubert DA, *et al.* 2012. *J. Exp Med.* 209:335. [PubMed](#)
 6. Dalli J, *et al.* 2012. *Blood.* 120:e60. [PubMed.](#)
 7. TrabANELLI S, *et al.* 2014. *J Immunol.* 192:1231. [PubMed](#)
 8. Herath S, *et al.* 2014. *PLoS One.* 9:88327. [PubMed](#)

Description: CD80, also known as B7-1, B7, and BB1, is a 60 kD single chain type I glycoprotein belonging to the immunoglobulin superfamily. CD80 is expressed on activated B and T cells, macrophages, and dendritic cells. CD80 binds to CD28 and CD152 (CTLA-4). Along with CD86, CD80 plays a critical role in regulation of T cell activation. The interaction of CD80 with CD28 provides a potent costimulatory signal for T cell activation through the CD3 complex, while its interaction with CTLA-4 provides an inhibitory signal for T cell activation.

Antigen 1. Freeman G, *et al.* 1991. *J. Exp. Med.* 174:625.

- References:**
2. Linsley P, *et al.* 1996. *Immunity* 4:535.
 3. Linsley P, *et al.* 1991. *J. Exp. Med.* 174:561.