

Biotin anti-human CD83

Catalog # / Size: 2126520 / 100 µg
2126515 / 25 µg

Clone: HB15e


Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



Monocytes-derived dendritic cells (induced with GM-CSF+IL-4+TNF-α) stained with biotinylated HB15e, followed by Sav-PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴.

Application References:

1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press New York.
2. Zhou L, *et al.* 1995. *J. Immunol.* 154:3821.
3. Cao W, *et al.* 2005. *Biochem. J.* 385:85.
4. Lore K, *et al.* 2002. *AIDS* 16:683. (IHC)
5. Cho H, *et al.* 2007. *Physiol Genomics* doi:10.1152/physiolgenomics.00051.2006

Description: CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member of the immunoglobulin superfamily, CD83 is expressed on a subset of dendritic cells, Langerhans cells, and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen presentation and/or lymphocyte activation, the precise function of this protein is unknown. CD83 is considered to be a useful marker for mature dendritic cells.

Antigen References:

1. Kozlow E, *et al.* 1993. *Blood* 81:454.
2. Zhou L, *et al.* 1992. *J. Immunol.* 149:735.
3. Zhou L, *et al.* 1995. *Blood* 86:3295.