

PE anti-human CD83

Catalog # / Size: 2126535 / 25 tests
2126540 / 100 tests

2126610 / 100 µg

Clone: HB15e


Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: test sizes: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
microg size: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: microg sizes: 0.2 mg/ml
test sizes: lot-specific



Monocyte-derived dendritic cells stained with HB15e PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴.

- Application References:**
1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press New York.
 2. Zhou L, *et al.* 1995. *J. Immunol.* 154:3821.
 3. Cao W, *et al.* 2005. *Biochem. J.* 385:85.
 4. Lore K, *et al.* 2002. *AIDS* 16:683. (IHC)
 5. Cho H, *et al.* 2007. *Physiol Genomics* doi:10.1152/physiolgenomics.00051.2006
 6. Belengio F, *et al.* 2013. *Immunobiology.* 218:76. [PubMed](#)
 7. Skrnjug I, *et al.* 2014. *PLoS One.* 9:95728. [PubMed](#)

Description: CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member of the immunoglobulin superfamily, CD83 is expressed on a subset of dendritic cells, Langerhans cells, and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen presentation and/or lymphocyte activation, the precise function of this protein is unknown. CD83 is considered to be a useful marker for mature dendritic cells.

- Antigen References:**
1. Kozlow E, *et al.* 1993. *Blood* 81:454.
 2. Zhou L, *et al.* 1992. *J. Immunol.* 149:735.
 3. Zhou L, *et al.* 1995. *Blood* 86:3295.

