

Brilliant Violet 421™ anti-human CD83

Catalog # / Size: 2126615 / 25 tests
2126620 / 100 tests

Clone: HB15e


Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human monocyte-derived dendritic cells (induced with GM-CSF+IL-4+TNF-α) were stained with CD83 (clone HB15e) Brilliant Violet 421™ (filled histogram) or mouse IgG1, κ Brilliant Violet 421™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴.

- Application References:**
1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press New York.
 2. Zhou L, *et al.* 1995. *J. Immunol.* 154:3821.
 3. Cao W, *et al.* 2005. *Biochem. J.* 385:85.
 4. Lore K, *et al.* 2002. *AIDS* 16:683. (IHC)
 5. Cho H, *et al.* 2007. *Physiol Genomics* doi:10.1152/physiolgenomics.00051.2006
 6. Royle CM, *et al.* 2014. *J Immunol.* 193:3538. [PubMed](#)

Description: CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member

of the immunoglobulin superfamily, CD83 is expressed on a subset of dendritic cells, Langerhans cells, and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen presentation and/or lymphocyte activation, the precise function of this protein is unknown. CD83 is considered to be a useful marker for mature dendritic cells.

**Antigen
References:**

1. Kozlow E, *et al.* 1993. *Blood* 81:454.
2. Zhou L, *et al.* 1992. *J. Immunol.* 149:735.
3. Zhou L, *et al.* 1995. *Blood* 86:3295.