

Alexa Fluor® 488 anti-human CD86

Catalog # / Size: 2127070 / 100 tests
2127065 / 25 tests

Clone: IT2.2

Isotype: Mouse IgG2b, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 488 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: VI CD86.8

Concentration: Lot-specific



Human peripheral blood monocytes with IT2.2 Alexa Fluor® 488

Applications:

- Applications:** Flow Cytometry
- Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
- * Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.
- Application Notes:** Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁶, Western blotting³, and blocking of T cell activation^{2,4,5}. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 305410).
- Application References:**
1. Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
 2. Dieu M. 1998. *J. Exp. Med.* 188:373. (Block)
 3. Esser M, *et al.* 2001. *J. Virol.* 75:6173. (WB)
 4. Jeannin P, *et al.* 1999. *J. Immunol.* 162:2044. (Block)
 5. Kapsogeorgou EK, *et al.* 2001. *J. Immunol.* 166:3107. (Block)
 6. Geissmann F, *et al.* 2001. *Blood* 97:1241. (IHC)

Description: CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, monocytes/macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is the ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86 binds to CD28 to transduce costimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can bind to CD152 as well, also known as CTLA-4, to deliver an inhibitory signal to T cells.

- Antigen** 1. Hathcock K, *et al.* 1996. *Adv. Immunol.* 62:131.
- References:** 2. June C, *et al.* 1994. *Immunol. Today* 15:321.