


Purified anti-human CD95 (Fas)


Catalog # / Size: 2128010 / 100 µg
Clone: DX2
Isotype: Mouse IgG1, κ
Immunogen: CD95 transfected L cells
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH7.2, containing 0.09% sodium azide.
Workshop Number: VI C-64
Concentration: 0.5



Human peripheral blood lymphocytes stained with DX2 PE/Cy5

Applications:

Applications: Flow Cytometry, Immunohistochemistry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



MCF7 breast cancer cells were stained with anti-human CD95 (clone DX2) using 1:100 dilution, followed by DyLight™ 649 anti-mouse Ig secondary antibody (red) plus DAPI staining for nuclei (blue). Cells were fixed with 4% PFA, permeabilized with 0.1%

Application Notes: The DX2 antibody is useful for inducing apoptosis of Fas-positive cells. Additional reported applications (for the relevant formats) include: *in vitro* induction of apoptosis³ (DX2 antibody is required to be cross-linked for effective induction of apoptosis) and immunohistochemical staining^{4,5} of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 305614).

Note: EOS9.1 antibody (Cat. No. 305704) can induce apoptosis without cross-linking.

Application References:

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. New York.
- Cifone M, *et al.* 1994. *J. Exp. Med.* 180:1547. (Apop)

4. Zietz C, *et al.* 2001. *Am. J. Pathol.* 159:963. (IHC)
 5. Sergi C, *et al.* 2000. *Am. J. Pathol.* 156:1589. (IHC)
 6. Xie S, *et al.* 2010. *J. Immunol.* 184:2289. (FC) [PubMed](#)
 7. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 8. Sestak K, *et al.* 2007. *Vet. Immunol. Immunopathol.* 119:21.
 9. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
 10. Dixit N, *et al.* 2012. *J. Immunol.* 189:5954. [PubMed](#)
-

Description: CD95 is a 45 kD single chain type I glycoprotein also known as Fas, APO-1, and TNFRSF6. It is a member of the TNF receptor superfamily. CD95 is expressed on T and B lymphocytes, monocytes, neutrophils, and fibroblasts. CD95 expression is upregulated by activation. The extracellular region of CD95 binds to CD178 (Fas ligand). CD178 binding to CD95 induces apoptosis and has been shown to play a role in the maintenance of peripheral tolerance.

Antigen 1. Krammer P, *et al.* 1994. *Immunol. Rev.* 142:175.
References: 2. Nagata S, *et al.* 1995. *Science* 267:1449.