

**PE/Fire™ 640 anti-human CD95 (Fas)**

**Catalog # / Size:** 2128290 / 100 tests  
2128285 / 25 tests

**Clone:** DX2

**Isotype:** Mouse IgG1, κ

**Immunogen:** CD95 transfected L cells


**Reactivity:** Human, Non-human primate

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Fire™ 640 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

**Workshop Number:** VI C-64

**Concentration:** Lot-specific



Human peripheral blood lymphocytes were stained with anti-human CD95 (Fas) (clone DX2) PE/Fire™ 640 (filled histogram), or mouse IgG1, κ PE/Fire™ 640 isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* PE/Fire™ 640 has a maximum excitation of 566 nm and a maximum emission of 639 nm.

**Application Notes:** The DX2 antibody is useful for inducing apoptosis of Fas-positive cells. Additional reported applications (for the relevant formats) include: *in vitro* induction of apoptosis<sup>3</sup> (DX2 antibody is required to be cross-linked for effective induction of apoptosis) and immunohistochemical staining<sup>4,5</sup> of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 305655 and 305656).

**Note:** EOS9.1 antibody can induce apoptosis without cross-linking.

**Application  
References:**

1. Schlossman S, et al. Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
  2. Kishimoto T, et al. Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. New York.
  3. Cifone M, et al. 1994. *J. Exp. Med.* 180:1547. (Apop)
  4. Zietz C, et al. 2001. *Am. J. Pathol.* 159:963. (IHC)
  5. Sergi C, et al. 2000. *Am. J. Pathol.* 156:1589. (IHC)
  6. Xie S, et al. 2010. *J. Immunol.* 184:2289. (FC) [PubMed](#)
  7. Yoshino N, et al. 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
  8. Sestak K, et al. 2007. *Vet. Immunol. Immunopathol.* 119:21.
  9. Rout N, et al. 2010. *PLoS One* 5:e9787. (FC)
  10. Dixit N, et al. 2012. *J. Immunol.* 189:5954. [PubMed](#)
- 

**Description:** CD95 is a 45 kD single chain type I glycoprotein also known as Fas, APO-1, and TNFRSF6. It is a member of the TNF receptor superfamily. CD95 is expressed on T and B lymphocytes, monocytes, neutrophils, and fibroblasts. CD95 expression is upregulated by activation. The extracellular region of CD95 binds to CD178 (Fas ligand). CD178 binding to CD95 induces apoptosis and has been shown to play a role in the maintenance of peripheral tolerance.

**Antigen  
References:**

1. Krammer P, et al. 1994. *Immunol Rev.* 142:175.
2. Nagata S, et al. 1995. *Science.* 267:1449.