

PE anti-human CD106

Catalog # / Size: 2129025 / 25 tests
2129030 / 100 tests

Clone: STA

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V A013

Concentration: Lot-specific



TNF- α stimulated HUVEC cells stained with STA PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence³, immunohistochemical staining of acetone-fixed frozen tissue sections, immunoprecipitation², and ELISA² capture for sCD106.

Application References:

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Leca G, *et al.* 1995. *J. Immunol.* 154:1069. (ELISA IP)
- Yen YT, *et al.* 2006. *J. Virol.* 80:2648. (IF) [PubMed](#)
- Wagner BJ, *et al.* 2011. *J Cell Sci.* 124:1644. [PubMed](#)
- Jin D, *et al.* 2011. *Regul Pept.* 166:21. [PubMed](#)
- Song K, *et al.* 2012. *Exp Cell Res.* 318:1707. [PubMed](#)
- Aomatsu E, *et al.* 2014. *Sci Rep.* 4:3652. [PubMed](#)

Description: CD106 is a 110 kD single chain type I glycoprotein also known as VCAM-1 and INCAM-110. It is expressed predominantly on activated vascular endothelium but has also been identified on follicular and interfollicular dendritic cells, some macrophages, bone marrow stromal cells, and non-vascular cell populations within joints, kidney, muscle, heart, placenta, and brain. Expression on endothelial cells as well as many other cells is induced by inflammatory stimuli and cytokines. Activated endothelial cells can release soluble forms of CD106 which can be detected in the blood. CD106 binds the integrins CD49d/CD29 (VLA-4) and $\alpha_4\beta_7$ that contribute to leukocyte adhesion, transmigration, and co-stimulation of T cell proliferation.

- Antigen** 1. Carlos T, *et al.* 1994. *Blood* 84:2068.
References: 2. Jones E, *et al.* 1995. *Nature* 373:539.