

FITC anti-human CD123

Catalog # / Size: 2130065 / 25 tests
2130070 / 100 tests

Clone: 6H6

Isotype: Mouse IgG1, κ


Immunogen: Human IL-3R α transfected COS cells.

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood leukocytes stained with CD45RO APC and 6H6 FITC (gated on lymphocyte population)

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone 6H6 does not inhibit IL-3 binding to low- or high-affinity IL-3Rs. Additional reported applications (for the relevant formats) include: Western blotting¹, immunoprecipitation¹, and immunohistochemical staining of acetone-fixed frozen sections².

- Application References:**
1. Sun Q, *et al.* 1996. *Blood* 87:83. (IP, WB)
 2. Herling M, *et al.* 2003. *Blood* 101:5007. (IHC)
 3. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
 4. Martin-Gayo E, *et al.* 2010. *Blood* 115:5366. [PubMed](#)
 5. Pritchard AL, *et al.* 2014. *PLoS One.* 9:106501. [PubMed](#)

Description: CD123 is the 70 kD transmembrane α chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common β chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the β chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.

Antigen References:

1. Miyajima A, *et al.* 1993. *Blood* 82:1960.