

APC/Fire™ 750 anti-human CD123

Catalog # / Size: 2130210 / 100 tests
2130205 / 25 tests

Clone: 6H6

Isotype: Mouse IgG1, κ

Immunogen: Human IL-3Rα transfected COS cells.


Reactivity: Human, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: 750 under optimal conditions.

Concentration: Lot-specific



Human lysed whole blood was stained with CD45RO FITC and CD123 (clone 6H6) APC/Fire™ 750 (left) or mouse IgG1, κ APC/Fire™ 750 isotype control (right).


Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: Clone 6H6 does not inhibit IL-3 binding to low- or high-affinity IL-3Rs. Additional reported applications (for the relevant formats) include: Western blotting¹, immunoprecipitation¹, and immunohistochemical staining of acetone-fixed frozen sections² and also paraformaldehyde fixed paraffin embedded tissue⁷.



- Application References:**
1. Sun Q, *et al.* 1996. *Blood* 87:83. (IP, WB)
 2. Herling M, *et al.* 2003. *Blood* 101:5007. (IHC)
 3. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
 4. Martin-Gayo E, *et al.* 2010. *Blood* 115:5366. [PubMed](#)
 5. Chen SC, *et al.* 2010. *Arch Dermatol Res.* 302:113. [PubMed](#)
 6. Liu Y, *et al.* 2012. *Food Chem Toxicol.* 50:1920. [PubMed](#)
 7. Peduzzi E, *et al.* 2007. *J. Invest. Dermatol.* 127:638. (IHC)

Description: CD123 is the 70 kD transmembrane α chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common β chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the β chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.

Antigen
References: 1. Miyajima A, *et al.* 1993. *Blood* 82:1960.