

**Alexa Fluor® 647 anti-human CD147**

**Catalog # / Size:** 2131045 / 25 tests  
2131050 / 100 tests

**Clone:** HIM6

**Isotype:** Mouse IgG1, κ

**Immunogen:** Human PBMCs


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Workshop Number:** VI N-L109

**Concentration:** Lot-specific




Human peripheral blood lymphocytes stained with HIM6 Alexa Fluor® 647

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. For immunofluorescence microscopy, a concentration range of 5-10 µg per ml is recommended. For immunohistochemical staining on formalin-fixed paraffin-embedded tissue sections, the suggested use of this reagent is 5.0 - 10 microg per ml. It is recommended that the reagent be titrated for optimal performance for each application.



HeLa cells were fixed with 1% paraformaldehyde (PFA) and then stained with 10 microg/ml of anti-human CD147 (clone HIM6) Alexa Fluor® 647 (red) for 3 hours at room temperature. Nuclei were counterstained with DAPI (blue). The image was captured by 2

\* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

**Application Notes:** Additional reported applications (for the relevant formats) include: inhibition of T cell activation<sup>2</sup>, immunohistochemical staining<sup>1,3</sup> of frozen tissue sections and formalin-fixed paraffin-embedded tissue sections, and Western blotting<sup>1</sup>. The LEAF™ Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 306206).

- Application** 1. Menashi S, *et al.* 2003. *Cancer Res.* 63:7575. (WB, IHC)  
**References:** 2. Woodhead VE, *et al.* 2000. *Int. Immunol.* 12:1051. (Block)  
3. Reimers N, *et al.* 2004. *Clin. Cancer Res.* 10:3422. (IHC)  
4. Porat-Shliom N, *et al.* 2013. *PLoS One.* 25:81897. [PubMed](#)
- 

**Description:** CD147, also known as neurothelin or basigin, is a member of the Ig superfamily. It is a 55-65 kD type I transmembrane glycoprotein which is primarily expressed on leukocytes, erythrocytes, platelets, and endothelial cells. CD147 is reported to have a function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain endothelia.

- Antigen**  
**References:** 1. Biswas C, *et al.* 1995. *Cancer Res.* 55:434.  
2. Fadool J, *et al.* 1993. *Dev. Dyn.* 196:252.  
3. Felzmann T, *et al.* 1991. *J. Clin. Immunol.* 11:205.