

PerCP/Cyanine5.5 anti-human CD147

Catalog # / Size: 2131095 / 25 tests
2131100 / 100 tests

Clone: HIM6

Isotype: Mouse IgG1, κ

Immunogen: Human PBMCs


Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cyanine5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: VI N-L109

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD147 (clone HIM6) PerCP/Cyanine5.5 (filled histogram) or mouse IgG1, κ PerCP/Cyanine5.5 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μl per million cells or 5 μl per 100 μl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of T cell activation², immunohistochemical staining^{1,3} of frozen tissue sections and formalin-fixed paraffin-embedded tissue sections, and Western blotting¹.

Application References:
 1. Biswas C, *et al.* 1995. *Cancer Res.* 55:434.
 2. Fadool J, *et al.* 1993. *Dev. Dyn.* 196:252.
 3. Felzmann T, *et al.* 1991. *J. Clin. Immunol.* 11:205.

Description: CD147, also known as neurothelin or basigin, is a member of the Ig superfamily. It is a 55-65 kD type I transmembrane glycoprotein which is primarily expressed on leukocytes, erythrocytes, platelets, and endothelial cells. CD147 is reported to have a function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain endothelia.

Antigen References:
 1. Biswas C, *et al.* 1995. *Cancer Res.* 55:434.
 2. Fadool J, *et al.* 1993. *Dev. Dyn.* 196:252.
 3. Felzmann T, *et al.* 1991. *J. Clin. Immunol.* 11:205.