

APC/Fire™ 750 anti-human TCR α/β

Catalog # / 2133680 / 100 tests
Size: 2133675 / 25 tests

Clone: IP26

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD01.01

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 FITC and anti-human TCRα/β (clone IP26) APC/Fire™ 750 (top) or mouse IgG1, κ APC/Fire™ 750 isotype control (bottom).

Applications:


Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: Additional reported applications (for the relevant formats) include: T cell activation. When co-staining with anti-CD3, we recommend using clone UCHT1, since we have confirmed that IP26 does not compete with this clone. Other anti-CD3 clones may compete out the binding of IP26.

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. (FC)
 - Joseph A, *et al.* 2008. *J. Virol.* 82:3078. (FC) [PubMed](#)
 - Pinto JP, *et al.* 2010. *Immunology.* 130:217. [PubMed](#)



Description: The IP26 antibody reacts with a monomorphic determinant of the α/β T-cell receptor, which is expressed on greater than 95% of normal peripheral blood CD3⁺ T cells. The α/β TCR recognizes a peptide bound to MHC leading to T-cell activation.

Antigen 1. Marchalonis J, et al. 2002. *J. Mol. Recognit.* 15:260.
References: