
APC/Fire[™] 810 anti-human TCR α/β

Catalog # / Size:	2133750 / 100 tests 2133745 / 25 tests
Clone:	IP26
lsotype:	Mouse IgG1, к
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 810 under optimal conditions.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)
Workshop Number:	V CD01.01
Concentration:	Lot-specific



Human peripheral blood lymphocytes were stained with anti-human CD3 FITC and antihuman TCR α/β (clone IP26) APC/Fire[™] 810 (left) or mouse IgG1, κ APC/Fire[™] 810 isotype control (right).

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
	* APC/Fire™ 810 has a maximum excitation of 650 nm and a maximum emission of 810 nm.
Application Notes:	Additional reported applications (for the relevant formats) include: T cell activation. When co-staining with anti-CD3, we recommend using clone UCHT1, since we have confirmed that IP26 does not compete with this clone. Other anti-CD3 clones may compete out the binding of IP26.
Application References:	 Schlossman S, et al. Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. (FC) Joseph A, et al. 2008. J. Virol. 82:3078. (FC) <u>PubMed</u> Pinto JP, et al. 2010. Immunology. 130:217. <u>PubMed</u>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Description:	The IP26 antibody reacts with a monomorphic determinant of the α/β T-cell receptor, which is expressed on greater than 95% of normal peripheral blood CD3 ⁺ T cells. The α/β TCR recognizes a peptide bound to MHC leading to T-cell activation.
Antigen	1. Marchalonis J, <i>et al.</i> 2002. <i>J. Mol. Recognit.</i> 15:260.

Antigen References: