SONY

Product Data Sheet

Brilliant Violet 570™ anti-human HLA-DR

Catalog # / Size: 2138185 / 25 tests
2138190 / 100 tests

Clone: L243

Isotype: Mouse IgG2a, κ

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 570™ under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with HLA-DR (clone L243) Brilliant Violet 570™ (filled histogram) or mouse IgG2a, κBrilliant Violet 570™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 570™ excites at 405 nm and emits at 570 nm. The bandpass filter 585/42 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 570™ is a trademark of Sirigen Group Ltd.

Application Notes: The L243 monoclonal antibody reacts with the HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. Clone L243 binds a conformational epitope on HLA-DRα which depends on the correct folding of the αβ heterodimer.[19]

Additional reported applications (for the relevant formats) include: immunoprecipitation[8], Western blotting[8], *in vitro* blocking of mixed lymphocyte reactions[9,10], depeletion of MHC class II cells[7], and immunohistochemical staining of acetone-fixed frozen sections[4,5]. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 307612). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 307648) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Application References:

1. Brodsky F. 1984. *Immunogenetics* 19:179.
2. Robbins P, *et al.* 1987. *Human Immunol.* 18:301.
3. Stites D, *et al.* 1986. *Clin. Immunol. Immunopathol.* 38:161.
4. Warnke R, *et al.* 1980. *J. Histochem. Cytochem.* 28:771. (IHC)
5. Engleman E, *et al.* 1981. *P. Natl. Acad. Sci. USA* 78:1791. (IHC)
6. Zipf T, *et al.* 1981. *Cancer Res.* 41:4786.
7. Goodier M, *et al.* 2000. *J. Immunol.* 165:139. (Depletion)
8. Esser M, *et al.* 2001. *J. Virol.* 75:6173. (IP, WB)
9. Kalka-Moll WM, *et al.* 2002. *J. Immunol.* 169:6149. (Block)
10. Wang RF, *et al.* 1999. *Science* 284:1351. (Block)
11. Zaba LC, *et al.* 2007. *J. Exp. Med.* 204:3183. PubMed
12. Fujita H, *et al.* 2009. *P. Natl. Acad. Sci. USA* 106:21795. PubMed
13. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) PubMed
14. Goncalves RM, *et al.* 2010. *Infect. Immun.* 78:4763. PubMed
15. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
16. Kim WK, *et al.* 2006. *Am. J. Pathol.* 168:822. (FC)
17. Stein R, *et al.* 2011. *Leuk. Lymphoma* 52:273.
18. Galkowska H, *et al.* 1996. *Vet. Immunol. Immunopathol.* 53:329.
19. Moro M, *et al.* 2005. *BMC Immunol.* 6:24.
20. Lauterbach N, *et al.* 2014. *Mol Immunol.* 59:19. PubMed

---

Description: HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to $CD4^+$ T cells.

Antigen References:

1. Levacher M, *et al.* 1990. *Clin. Exp. Immunol.* 81:177.
2. Terstappen L, *et al.* 1990. *J. Leukocyte Biol.* 48:138.
3. Edwards JA, *et al.* 1986. *J. Immunol.* 137:490.
4. van Es A, *et al.* 1984. *Transplantation* 37:65.
5. O'Doherty U, *et al.* 1994. *Immunology* 82:487.
6. Thomas R, *et al.* 1994. *J. Immunol.* 153:4016.
7. Grouard G, *et al.* 1996. *Nature* 384:364.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.