


## Brilliant Violet 510™ anti-human Perforin


<b>Catalog # / Size:</b>	2140595 / 25 tests 2140600 / 100 tests
<b>Clone:</b>	dG9
<b>Isotype:</b>	Mouse IgG2b, $\kappa$
<b>Immunogen:</b>	Purified granules from the human lymphoma cell line
<b>Reactivity:</b>	Human
<b>Preparation:</b>	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody.
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
<b>Concentration:</b>	Lot-specific



Human peripheral blood lymphocytes were surface stained with CD8 APC and then intracellularly stained with perforin (clone dG9) Brilliant Violet 510™ (top), or mouse IgG2b,  $\kappa$  Brilliant Violet 510™ isotype control (bottom).

## Applications:

<b>Applications:</b>	Flow Cytometry
<b>Recommended Usage:</b>	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 5$ microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the

buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** Clone dG9 primarily recognizes perforin associated with cytotoxic granules<sup>9</sup>. Additional reported applications (for the relevant formats) include: immunoprecipitation, intracellular flow cytometric analysis and immunofluorescence microscopy<sup>5,7</sup>, and immunohistochemical staining of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections<sup>1,4</sup>.

Does not cross-react with mouse<sup>>1</sup>.</sup>

**Application References:**

1. Hameed A, *et al.* 1992. *Am. J. Pathol.* 140:1025. (IHC)
2. Schaerli P, *et al.* 2004. *J. Exp. Med.* 199:1265.
3. Watanabe N, *et al.* 1997. *Blood* 90:3662.
4. Mauad T, *et al.* 2004. *Pediatr. Pulmonol.* 38:233. (IHC)
5. Barrat FJ, *et al.* 1999. *P. Natl. Acad. Sci. USA* 96:8645. (IF)
6. Chen H, *et al.* 2005. *J. Immunol.* 175:591.
7. Bryceson YT, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-074468. (IF)
8. Wood SM, *et al.* 2009. *Blood* 114:4117. [PubMed](#)
9. Makedonas G, *et al.* 2010. *PLoS Pathog.* 6:e1000798.

---

**Description:** Perforin is a 70 kD cytolytic protein that is expressed in the cytoplasmic granules of cytotoxic T lymphocytes (CTLs) and natural killer (NK) cells. Perforin is one of the major effector molecules used by cytotoxic T cells and NK cells to mediate targeted cell lysis.

**Antigen References:**

1. Lieberman J. 2003. *Nat. Rev. Immunol.* 3:361.
2. Trapani J, *et al.* 2002. *Nat. Rev. Immunol.* 2:735.