

**APC anti-human CD275 (B7-H2, ICOSL)**

**Catalog # / Size:** 2147040 / 100 tests  
2147035 / 25 tests

**Clone:** 2D3

**Isotype:** Mouse IgG2b,  $\kappa$


**Immunogen:** Human B7H2-mIg fusion protein

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



Human Burkitt's lymphoma cell line Daudi was stained with CD275 (clone 2D3) APC (filled histogram) or Mouse IgG2b,  $\kappa$  APC isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:** 1. Kurosawa S, *et al.* 2003. *Am. J. Respir. Cell Mol. Biol.* 28:563.

**Description:** B7-H2, a member of the B7 family and the immunoglobulin superfamily, is a 40 kD protein also known as B7RP-1, B7h, B7-H2, GL50 and ICOS Ligand. Human B7-H2 is expressed by activated monocytes/macrophages and dendritic cells. B7-H2 binds to a CD28-like receptor, inducible costimulator molecule (ICOS, AILIM, CRP-1), which is expressed by activated T cells. The interaction of ICOS with B7-H2 plays an important role in the T cell costimulation pathway.

**Antigen References:** 1. Wang S, *et al.* 2002. *J. Biol. Chem.* 96:2808.  
2. Wong SC, *et al.* 2003. *Blood* 102:1831.