


Alexa Fluor® 647 anti-human CD137 (4-1BB)

| | |
|--------------------------|---|
| Catalog # / Size: | 2149120 / 100 tests 2149115 / 25 tests |
| Clone: | 4B4-1 |
| Isotype: | Mouse IgG1, κ |
| Immunogen: | Ectodomain of recombinant human 4-1BB fusion protein |
| Reactivity: | Human |
| Preparation: | The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions. |
| Formulation: | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA). |
| Workshop Number: | VI C-7 |
| Concentration: | Lot-specific |



PHA-stimulated (three days) human peripheral blood lymphocytes were stained with CD137 (clone 4B4-1) Alexa Fluor® 647 (filled histogram) or mouse IgG1, κ Alexa Fluor® 647 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,4}, inhibition of cytokine production^{2,3}, and ELISA. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 309804) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated anti-mouse IgG second step (Cat. No. 405303), followed by Streptavidin-PE (Cat. No. 405204)).

Application References:

1. Garni-Wagner B, *et al.* 1996. *Cell. Immunol.* 169:91. (IP)
2. Salih HR, *et al.* 2000. *J. Immunol.* 165:2903. (FA)
3. Kienzle G, *et al.* 2000. *Int. Immunol.* 12:73. (FA)
4. Langstein J, *et al.* 1998. *J. Immunol.* 160:2488. (IP)

Description: CDw137 is a 39 kD transmembrane protein also known as 4-1BB. It is expressed on activated T cells. CDw137 is a type I membrane protein and a member of the tumor necrosis factor receptor superfamily. CDw137 appears to be important for T cell proliferation and survival, and induces monocyte activation through its interaction with 4-1BB ligand.

Antigen References:

1. Gruss H, *et al.* 1995. *Blood* 85:3378.
2. Sica G, *et al.* 2000. *Adv. Exp. Med. Biol.* 465:355.

3. Alderson M, *et al.* 1994. *Eur. J. Immunol.* 24:2219.
4. Schwarz H, *et al.* 199