

Brilliant Violet 711™ anti-human CD137 (4-1BB)

Catalog # / Size:	2149155 / 25 tests 2149160 / 100 tests
Clone:	4B4-1
Isotype:	Mouse IgG1, κ
Immunogen:	Ectodomain of recombinant human 4-1BB fusion protein
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 711™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 711™ and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Workshop Number:	VI C-7
Concentration:	0.2

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 711™ excites at 405 nm and emits at 711 nm. The bandpass filter 710/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 711™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes:	Additional reported applications (for the relevant formats) include: immunoprecipitation ^{1,4} , inhibition of cytokine production ^{2,3} , and ELISA. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 309804) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated anti-mouse IgG second step (Cat. No. 405303), followed by Streptavidin-PE (Cat. No. 405204)).
---------------------------	--

- Application** 1. Garni-Wagner B, *et al.* 1996. *Cell. Immunol.* 169:91. (IP)
- References:** 2. Salih HR, *et al.* 2000. *J. Immunol.* 165:2903. (FA)
3. Kienzle G, *et al.* 2000. *Int. Immunol.* 12:73. (FA)
4. Langstein J, *et al.* 1998. *J. Immunol.* 160:2488. (IP)
-

Description: CD137 is a 39 kD transmembrane protein also known as 4-1BB. It is expressed on activated T cells. CD137 is a type I membrane protein and a member of the tumor necrosis factor receptor superfamily. CD137 appears to be important for T cell proliferation and survival, and induces monocyte activation through its interaction with 4-1BB ligand.

- Antigen** 1. Gruss H, *et al.* 1995. *Blood* 85:3378.
- References:** 2. Sica G, *et al.* 2000. *Adv. Exp. Med. Biol.* 465:355.
3. Alderson M, *et al.* 1994. *Eur. J. Immunol.* 24:2219.
4. Schwarz H, *et al.* 199